## List of new drugs approved in the year 2019 till date

S.No	Name of drug	Indication	Date of issue
1	Fenspiride hydrochloride film coated extended release tablet 80 mg and Fenspiride hydrochloride bulk	Acute Rhinosinusitis     Moderate persistent asthma as an addon therapy	04.02.2019
2	Bilastine tablets 20 mg and Bilastine Bulk	For symptomatic treatment of allergic rhino-conjuctivitis (seasonal and perennial) and utricaria in adults	06.02.2019
3	Iguratimod film coated tablets 25mg and Iguratimod Bulk	For the treatment of active rheumatoid arthritis symptoms	18.02.2019
4	Fingolimod Capsules 0.5mg and Fingolimod hydrochloridebulk	For the treatment of patients with relapsing forms of multiple sclerosis (MS) to reduce the frequency of clinical exacerbations and to delay the accumulation of physical disability.	25.03.2019
5	Remogliflozin etabonate bulk and Remogliflozin etabonate film coated tablets 100 mg	Indicated in adults aged 18 years and older with type2 diabetes mellitus to improve glycemic control as:-  • Monotherapy when diet and exercise alone do not provide adequate glyceamic control.  • Add on therapy with metformin, together with diet and exercise, when these do not provide adequate glyceamic control.	26.04.2019
6	Concentrate of Proteolytic enzyme enriched in Bromelain topical gel	Indicated for removal of eschar in adults with deep partial-and full-thickness thermal burns	17.05.2019
7	Menotrophin injection 600 IU/ml, 1 ml & 2 ml multi dose vial (Highly purified)	Menotropin is indicated for the treatment of female and male infertility in the following conditions:  Anovulation, including Polycystic Ovarian Disease (PCOD), in women who have been unresponsive to the treatment with	17.06.2019

		Clomiphene citrate.	
8	Rifapentine 150mg film coated tablet	Clomiphene citrate.  Women undegoing controlled ovarian hyperstimulation to induce the development of multiple follicles for Assisted Reproductive Technologies (ART).  Hypogonadotropic hypogonadism in men.  "For the treatment of latent tuberculosis infection caused by Mycobacterium tuberculosis in adults and children 2 years and older who are at high risk of progression to tuberculosis disease (including those in close contact with active tuberculosis patients, recent conversion to a positive tuberculin skin test, HIV-infected patients, or those with pulmonary fibrosis on radiograph).  Active tuberculosis disease should be ruled out before initiating treatment for latent tuberculosis infection.  Rifapentine Tablets 150mg must always be used in combination with isoniazid as a 12-week once-weekly regimen for the treatment of latent tuberculosis infection".	08.07.2019
9	FDC of Bictegravir, Emtricitabine and Tenofovir alafenamide 50mg/200mg/25mg tablets Levomilnacipran ER capsules 20mg/40mg/80mg/120mg	For the treatment of human immunodeficiency virus type 1 (HIV-1) infection in adults who have no antiretroviral treatment history or to replace the current antiretroviral regimen in those who are virologically suppressed (HIV-1 RNA less than 50 copies per ml) on a stable antiretroviral regimen for at least 3 month with no history of treatment failure and no known substitutions associated with resistance to Bictegravir, Emtricitabine and Tenofovir alafenamide.	
10	zumg/4umg/8umg/12umg		13.00.2019
11	Bosutinib bulk and Bosutinib tablets 100mg,	For the treatmnet of adult patients with:  1) Newly diagnosed chronic phase (CP)	09.09.2019

	400mg, 500mg	Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+CML) 2) Chronic phase, accelerated phase (AP), or blast phase (BP) Ph+CML with resistance or intolerance to prior therapy.	
12	Algard - R-82 (octyldecyl dimethyl ammonium chloride 6.510 w/w + Dioctyl dimethyl ammonium chloride 2.604% w/w + Didecyl dimethyl ammonium chloride 3.906% w/w + Alkyl dimethyl benzyl ammonium chloride 8.680% w/w	Surface disinfectant for persistent disinfection of all inanimate surfaces (Against pathogens as annexed)	09.09.2019
13	Algard HWS-256 (didecyl dimethyl ammonium chloride 10.14% w/w + n-Alkyl dimethyl benzyl ammonium chloride 6.76% w/w	Surface disinfectant for persistent disinfection of all inanimate surfaces (Against pathogens as annexed)	09.09.2019
14	Safinamide methane sulphonate bulk drug and Safinamide tablets 50, 100mg	For the treatment of adult patients with idiopathic Parkinson's disease (PD) as addon therapy to a stable dose of Levodopa (Ldopa) alone or in combination with other Parkinson's disease (PD) medicinal products in mid to late stage fluctuating patients.	09.09.2019
15	Droxidopa bulk drug and Droxidopa capsules 200mg/300mg	For the treatment of orthostatic dizziness, light headedness or the "feeling that you are about to black out" in adult patients with symptomatic neurogenic orthostatic hypotension caused by primary autonomic failure (Parkinson's Disease multiple system atrophy (MSA) and pure autonomic failure), dopamine beta-hydroxylase deficiency and non-diabetic autonomic neuropathy.	24.09.2019
16	Acalabrutinib 100mg capsules	For the treatment of patients with mantle cell lymphoma (MCL) who have received atleast one prior therapy	18.10.2019
	Abemaciclib 50mg, 100mg, 150mg and 200mg film coated tablets	(i) Abemaciclib is indicated for the treatment of women with hormone receptor (HR) positive, human epidermal growth factor receptor 2 (HER2) negative locally advanced or metastatic breast cancer in combination with an aromatase	18.11.2019

		inhibitor or fulvestrant as initial endocrine-	l l
		based therapy, or in women who have	
		received prior endocrine therapy.	
		• In pre- or perimenopausal women, the	
		endocrine therapy should be combined	
		with a luteinising hormone- releasing	
		hormone (LHRH) agonist.	
		(ii) As monotherapy for the treatment of	
		adult patients with HR-positive, HER2-	
		negative advanced or metastatic breast	
		cancer with disease progression following	
		endocrine therapy and prior chemotherapy	
		in the metastatic setting.	
		in the metastatic setting.	
	S: 1911 1 11 11 1 1 1 1 1 1 1 1 1 1 1 1 1	To treat glaucoma and ocular hypertension	
	Ripasudil hydrochloride hydrate bulk drug and Ripasudil Eye drops 0.4% w/v	when other medicines for glaucoma have	20.11.2019
18		insufficient effect or cannot be used.	
	Dinaraya shlaria asid cancentrate and	Indicated for wound healing in diabetic	_
	Diperoxochloric acid concentrate and	neuropathic ulcers of skin and	20.11.2019
19	Diperoxochloric acid topical solution	subcutaneous tissues reduction	20.11.2019
	Endoxifen citrate bulk and Endoxifen tablets	For the acute treatment of manic episodes	
20	8 mg	with or without mixed features of Bipolar I	10.12.2019
		disorder	10.12.2013