



सत्यमेव जयते

**GOVERNMENT OF INDIA**  
**CENTRAL DRUGS STANDARD CONTROL**  
**ORGANISATION (Headquarter)**  
(Directorate General of Health Services)  
Ministry of Health & Family Welfare  
FDA Bhawan, Kotla Road,  
New Delhi – 110002  
Phone No.: 91-11-23216367  
Fax No.: 91-11-23236973  
E-Mail : dci@nic.in

File No. CT/22/000083

To

M/s IR Innovate Research Private Limited,  
C 1231, Rajajipuram, Lucknow–226017, Uttar Pradesh, India.  
Tel. No.: 9971169602; Fax: 9971169602  
E-mail: REGULATORY@INNOVATE-RESEARCH.COM

Sir,

With reference to your application vide SUGAM F. no. GCT/CT04/FF/2022/33472 (GCT/83/22) dated 12-AUG-2022, please find enclosed herewith the permission in Form CT-06 for conduct of phase III clinical trial titled, “ **A Randomized, Double-Blind Clinical Study of the Efficacy and Safety of BCD-201 (JSC BIOCAD) and Keytruda® in Patients with Unresectable or Metastatic Melanoma**” vide **Protocol No.: BCD-201-2, Amendment 1, Version 2.0, dated 14-JUL-2022 and India-Addendum Version 1.0 dated 16-NOV-2022** with a total of up to 50 subjects under the provisions of New Drugs and Clinical Trial Rules, 2019.

The permission granted by the Central Licensing Authority to conduct clinical trial shall be subject to following conditions, specifically that: -

- 1) **the firm should submit the Interim Study Report no. 1 of approx. 89 subjects for review of the SEC/CDSCO and further continuation of the trial in the country;**
- 2) **SAEs including death irrespective of its cause i.e. PD should be reported to the CDSCO as per provisions of the New Drugs and Clinical Trial Rules, 2019 during the trial;**
- 3) Clinical trial at each site shall be initiated after approval of the clinical trial protocol and other related documents by the Institutional Ethics Committee of that site, registered with the Central Licencing Authority under Rule 8;
- 4) where a clinical trial site does not have its own Ethics Committee, clinical trial at that site may be initiated after obtaining approval of the protocol from the Ethics Committee of another trial site; or an independent Ethics Committee for clinical trial constituted in accordance with the provisions of Rule 7:

Provided that the approving Ethics Committee for clinical trial shall in such case be responsible for the study at the trial site or the centre, as the case may be:

Provided further that the approving Ethics Committee and the clinical trial site or the bioavailability and bioequivalence centre, as the case may be, shall be located within the same city or within a radius of 50 kms of the clinical trial site;

- 5)** in case an ethics committee of a clinical trial site rejects the approval of the protocol, the details of the same shall be submitted to the Central Licensing Authority prior to seeking approval of another Ethics Committee for the protocol for conduct of the clinical trial at the same site;
- 6)** the Central Licencing Authority shall be informed about the approval granted by the Ethics Committee within a period of fifteen working days of the grant of such approval;
- 7)** clinical trial shall be registered with the Clinical Trial Registry of India maintained by the Indian Council of Medical Research before enrolling the first subject for the trial;
- 8)** clinical trial shall be conducted in accordance with the approved clinical trial protocol and other related documents and as per requirements of Good Clinical Practices Guidelines and the provisions of these rules;
- 9)** status of enrolment of the trial subjects shall be submitted to the Central Licencing Authority on quarterly basis or as appropriate as per the duration of treatment in accordance with the approved clinical trial protocol, whichever is earlier;
- 10)** six monthly status report of each clinical trial, as to whether it is ongoing, completed or terminated, shall be submitted to the Central Licencing Authority electronically in the SUGAM portal;
- 11)** in case of termination of any clinical trial the detailed reasons for such termination shall be communicated to the Central Licencing Authority within thirty working days of such termination;
- 12)** any report of serious adverse event occurring during clinical trial to a subject of clinical trial, shall, after due analysis, be forwarded to the Central Licencing Authority, the chairperson of the Ethics Committee and the institute where the trial has been conducted within fourteen days of its occurrence as per Table 5 of the Third Schedule and in compliance with the procedures as specified in Chapter VI;
- 13)** in case of injury during clinical trial to the subject of such trial, complete medical management and compensation shall be provided in accordance with Chapter VI and details of compensation provided in such cases shall be intimated to the Central Licencing Authority within thirty working days of the receipt of order issued by Central Licencing Authority in accordance with the provisions of the said Chapter;
- 14)** in case of clinical trial related death or permanent disability of any subject of such trial during the trial, compensation shall be provided in accordance with Chapter VI and details of compensation provided in such cases shall be intimated to the Central Licencing Authority within thirty working days of receipt of the order issued by the Central Licencing Authority in accordance with the provisions of the said Chapter;
- 15)** the premises of the sponsor including his representatives and clinical trial sites, shall be open for inspection by officers of the Central Licencing Authority who may be accompanied by officers of the State Licencing Authority or outside experts as authorised by the Central Licencing Authority, to verify compliance of the requirements of these rules and Good Clinical Practices Guidelines, to inspect, search and seize any record, result, document, investigational product, related to clinical trial and furnish reply to query raised by the said officer in relation to clinical trial;

- 16) where the new drug or investigational new drug is found to be useful in clinical development, the sponsor shall submit an application to the Central Licencing Authority for permission to import or manufacture for sale or for distribution of new drug in India, in accordance with Chapter X of these rules, unless otherwise justified;
- 17) the laboratory owned by any person or a company or any other legal entity and utilised by that person to whom permission for clinical trial has been granted used for research and development, shall be deemed to be registered with the Central Licensing Authority and may be used for test or analysis of any drug for and on behalf of Central Licensing Authority;
- 18) the Central Licencing Authority may, if considered necessary, impose any other condition in writing with justification, in respect of specific clinical trials, regarding the objective, design, subject population, subject eligibility, assessment, conduct and treatment of such specific clinical trial;
- 19) the sponsor and the investigator shall maintain the data integrity of the data generated during clinical trial.
- 20) merely granting permission to conduct the clinical trial with the Investigational Drug Product does not convey or imply that, based on the clinical trial study data generated with the investigational drug, permission to market this drug in the country will automatically be granted to you;
- 21) the permission to initiate clinical trial granted under rule 22 in form CT-06 or automatic approval under rule 23 in Form CT 4A shall remain valid for a period of **two years** from the date of its issue, unless extended by the Central Licencing Authority.

Yours faithfully,

VENUG  
OPAL G  
SOMANI  
(Dr. V. G. Somani)

Digitally signed by VENUGOPAL G SOMANI  
DN: c=IN, o=CENTRAL DRUGS STANDARD CONTROL ORGANIZATION, ou=DRUGS CONTROLLER GENERAL (INDIA), pseudonym=112046a34e21727bach255748732c456d249073e55e953d9cb5b58748ee3113c, postalCode=110002, st=DELHI, serialNumber=e97f241aa0c0bba41522525ffca0705074b4997e6b2f4b5c8d81cf282adeaf5d, cn=VENUGOPAL G SOMANI

Drugs Controller General (India) &  
Central Licencing Authority

**FORM CT-06**

(See rules 22,25,26,29 and 30)

**PERMISSION TO CONDUCT CLINICAL TRIAL OF NEW DRUG OR INVESTIGATIONAL NEW DRUG**

1. The Central Licensing Authority hereby permits **M/s IR Innovate Research Private Limited, C 1231, Rajajipuram, Lucknow-226017, Uttar Pradesh, India, Tel. No.: 9971169602; Fax: 9971169602 , E-mail: REGULATORY@INNOVATE-RESEARCH.COM** to conduct clinical trial of the new drug or investigational new drug as per **Protocol No.: BCD-201-2, Amendment 1, Version 2.0, dated 14-JUL-2022 and India-Addendum Version 1.0 dated 16-NOV-2022** in the below mentioned clinical trial sites [As per Annexure]. -
2. Details of new drug or investigational new drug and clinical trial site [As per Annexure].
3. This permission is subject to the conditions prescribed in part A of Chapter V of the New Drugs and Clinical Trials Rules, 2019 under the Drugs and Cosmetics Act, 1940.

Place: New Delhi

Date \_\_\_\_\_

VENUGO

PAL G

SOMANI

(Dr. V. G. Somani)

Drugs Controller General (India) &  
Central Licensing Authority

**Note:** The permission to initiate clinical trial granted under rule 22 in form CT-06 shall remain valid for a period of **two years** from the date of its issue, unless extended by the Central Licencing Authority.

**Annexure:**

Details of new drug or investigational new drug:

|                                                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |                          |          |  |                                    |  |  |  |         |         |                |        |                     |         |          |          |         |        |          |          |               |        |          |          |                                     |          |          |          |           |          |          |          |                      |              |                          |          |
|---------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------|----------|--|------------------------------------|--|--|--|---------|---------|----------------|--------|---------------------|---------|----------|----------|---------|--------|----------|----------|---------------|--------|----------|----------|-------------------------------------|----------|----------|----------|-----------|----------|----------|----------|----------------------|--------------|--------------------------|----------|
| Names of the new drug or investigational new drug | BCD-201                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                          |          |  |                                    |  |  |  |         |         |                |        |                     |         |          |          |         |        |          |          |               |        |          |          |                                     |          |          |          |           |          |          |          |                      |              |                          |          |
| Therapeutic class:                                | Anti-PD-1 monoclonal antibody                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |                          |          |  |                                    |  |  |  |         |         |                |        |                     |         |          |          |         |        |          |          |               |        |          |          |                                     |          |          |          |           |          |          |          |                      |              |                          |          |
| Dosage form:                                      | Concentrate solution for infusion                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |                          |          |  |                                    |  |  |  |         |         |                |        |                     |         |          |          |         |        |          |          |               |        |          |          |                                     |          |          |          |           |          |          |          |                      |              |                          |          |
| Composition:                                      | <table><tr><td colspan="4">BCD-201 DP composition (per 1 mL):</td></tr><tr><td>BCD-201</td><td>25.0 mg</td><td>In-house spec.</td><td>Active</td></tr><tr><td>Trehalose dihydrate</td><td>80.0 mg</td><td>Ph. Eur.</td><td>Inactive</td></tr><tr><td>Glycine</td><td>1.5 mg</td><td>Ph. Eur.</td><td>Inactive</td></tr><tr><td>Poloxamer 188</td><td>1.0 mg</td><td>Ph. Eur.</td><td>Inactive</td></tr><tr><td>Histidine hydrochloride monohydrate</td><td>0.750 mg</td><td>Ph. Eur.</td><td>Inactive</td></tr><tr><td>Histidine</td><td>0.221 mg</td><td>Ph. Eur.</td><td>Inactive</td></tr><tr><td>Water for injections</td><td>Up to 1.0 mL</td><td>Ph. Eur./ In house spec.</td><td>Inactive</td></tr></table> |                          |          |  | BCD-201 DP composition (per 1 mL): |  |  |  | BCD-201 | 25.0 mg | In-house spec. | Active | Trehalose dihydrate | 80.0 mg | Ph. Eur. | Inactive | Glycine | 1.5 mg | Ph. Eur. | Inactive | Poloxamer 188 | 1.0 mg | Ph. Eur. | Inactive | Histidine hydrochloride monohydrate | 0.750 mg | Ph. Eur. | Inactive | Histidine | 0.221 mg | Ph. Eur. | Inactive | Water for injections | Up to 1.0 mL | Ph. Eur./ In house spec. | Inactive |
| BCD-201 DP composition (per 1 mL):                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |                          |          |  |                                    |  |  |  |         |         |                |        |                     |         |          |          |         |        |          |          |               |        |          |          |                                     |          |          |          |           |          |          |          |                      |              |                          |          |
| BCD-201                                           | 25.0 mg                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | In-house spec.           | Active   |  |                                    |  |  |  |         |         |                |        |                     |         |          |          |         |        |          |          |               |        |          |          |                                     |          |          |          |           |          |          |          |                      |              |                          |          |
| Trehalose dihydrate                               | 80.0 mg                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | Ph. Eur.                 | Inactive |  |                                    |  |  |  |         |         |                |        |                     |         |          |          |         |        |          |          |               |        |          |          |                                     |          |          |          |           |          |          |          |                      |              |                          |          |
| Glycine                                           | 1.5 mg                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | Ph. Eur.                 | Inactive |  |                                    |  |  |  |         |         |                |        |                     |         |          |          |         |        |          |          |               |        |          |          |                                     |          |          |          |           |          |          |          |                      |              |                          |          |
| Poloxamer 188                                     | 1.0 mg                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | Ph. Eur.                 | Inactive |  |                                    |  |  |  |         |         |                |        |                     |         |          |          |         |        |          |          |               |        |          |          |                                     |          |          |          |           |          |          |          |                      |              |                          |          |
| Histidine hydrochloride monohydrate               | 0.750 mg                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | Ph. Eur.                 | Inactive |  |                                    |  |  |  |         |         |                |        |                     |         |          |          |         |        |          |          |               |        |          |          |                                     |          |          |          |           |          |          |          |                      |              |                          |          |
| Histidine                                         | 0.221 mg                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | Ph. Eur.                 | Inactive |  |                                    |  |  |  |         |         |                |        |                     |         |          |          |         |        |          |          |               |        |          |          |                                     |          |          |          |           |          |          |          |                      |              |                          |          |
| Water for injections                              | Up to 1.0 mL                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | Ph. Eur./ In house spec. | Inactive |  |                                    |  |  |  |         |         |                |        |                     |         |          |          |         |        |          |          |               |        |          |          |                                     |          |          |          |           |          |          |          |                      |              |                          |          |
| Indication:                                       | Unresectable or Metastatic Melanoma                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |                          |          |  |                                    |  |  |  |         |         |                |        |                     |         |          |          |         |        |          |          |               |        |          |          |                                     |          |          |          |           |          |          |          |                      |              |                          |          |

**Annexure:**

Details of clinical trial site:

| <b>S.No.</b> | <b>Names and address of clinical trial site</b>                                                                                                                                                                                                                                                                                 | <b>Ethics committee details</b>                                                                                                                                                                                 | <b>Name of Investigator</b> |
|--------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------|
| <b>1.</b>    | Unique Hospital<br>Multispeciality and Research<br>Institute, Opp. Kiran Motor,<br>Nr. Canal, Civil Hospital<br>Char Rasta, Sosyo Circle<br>Lane, off Ring Road, Surt-<br>395002, Gujarat<br><br>Unique Hospital Corporate<br>Office, B-9 Shivkrupa<br>Society, Ambanagar, Behind<br>Unique Hospital, Surat-<br>395002, Gujarat | Ethics Committee Unique Hospital<br>Multispeciality and Research<br>Institute, Opp. Kiran Motor, Nr.<br>Canal, Civil Hospital Char Rasta,<br>Sosyo Circle Lane, off Ring Road,<br>Surat-395002, Gujarat         | Dr. Tanveer M<br>Maksud     |
| <b>2.</b>    | Somani Hospital, Oncology<br>Department, Somani<br>Hospital, 277, Shri Gopal<br>Nagar, 80 Ft. Road,<br>Gopalpura Bypass, Jaipur-<br>302019, Rajasthan                                                                                                                                                                           | Somani Hospital Ethics<br>Committee, Somani Hospital, 277,<br>Shri Gopal Nagar, 80 Ft. Road,<br>Gopalpura Bypass, Jaipur-<br>302019, Rajasthan                                                                  | Dr. Naresh<br>Somani        |
| <b>3.</b>    | Marathwada Cancer Center.<br>Aurangabad, Near Jama<br>masjid. Opp humkhas<br>Maidan Marathwada<br>Regional Cancer Center and<br>Research Institute<br>Aurangabad-431001,M.S                                                                                                                                                     | Institutional Ethics Committee<br>Government Medical College<br>Aurangabad, Department of<br>Pharmacology Government<br>Medical College, Aurangabad-<br>431001, M.S                                             | Dr. Jitendra<br>Patel       |
| <b>4.</b>    | Grant Medical Foundation<br>Ruby Hall Clinic, 40,<br>Sassoon Road, Pune,<br>Maharashtra-411001                                                                                                                                                                                                                                  | Institutional Ethics Committee<br>Poona Medical Foundation, E-4 C<br>to E-4 F, 4 <sup>th</sup> Floor, Fifth Avenue<br>Condominium, Dhole Patil Road,<br>Pune, Maharashtra-411001                                | Dr. Jain Minish<br>Mahendra |
| <b>5.</b>    | Basavatarakam Indo<br>American Cancer Hospital &<br>Research Institute, Road<br>No. 10, IAS Officers<br>Quarters, Nandi Nagar,<br>Banjara Hills, Hyderabad,<br>Telangana-500034                                                                                                                                                 | Institutional Ethics Committee,<br>Basavatarakam Indo American<br>Cancer Hospital & Research<br>Institute, Road No. 10, IAS<br>Officers Quarters, Nandi Nagar,<br>Banjara Hills, Hyderabad,<br>Telangana-500034 | Dr. Ayyagari<br>Santa       |
| <b>6.</b>    | Jawaharlal Institute of<br>Postgraduate Medical<br>Education & research<br>(JIPMER), Department of<br>Oncology, Dhanvantari<br>Nagar, Gorimedu,<br>Puducherry-605006                                                                                                                                                            | IEC Intervention Studies, JIPMER,<br>Dhanvantari Nagar, Puducherry-<br>605006                                                                                                                                   | Dr. Smita Kayal             |

\*\*\*\*\*