

**SUMMARY OF PRODUCT CHARACTERISTICS**

**Doc. No. SPC/71108 Ver.3**

**1. NAME OF THE MEDICINAL PRODUCT.**

- Novel Corona Virus 2019-nCoV Vaccine (Recombinant)

**2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

**Each 0.1ml contains:**

DNA plasmid construct with spike protein gene region from SARS- 1.0 mg  
CoV-2 virus produced in *E.coli*  
Phosphate Buffered saline q.s.

**3. PHARMACEUTICAL FORM**

Solution for Intradermal Injection.

Each dose consists of three shots of 0.1 mL each

<b>CAUTION – Dose and Regimen Selection</b>	
3mg – 2 Dose Regimen: 3 shots of 0.1 ml each should be given on day 0 and 28	
<i>3x 0.1ml</i> ●	<i>3x 0.1ml</i> ●
<i>Day 0</i>	<i>Day 28</i>
It is recommended to use the vaccine only with Pharmajet Device. Using it with conventional needle and syringe will not lead to optimal immunogenicity response and will affect the efficacy of the vaccine.	

**4. CLINICAL PARTICULARS**

**4.1 Therapeutic indications**

ZYCOV-D® is indicated for active immunisation to prevent COVID-19 caused by SARSCoV-2 in individuals 12 years of age and older when given in two separate doses of 3mg (0.3ml) each to be given at an interval of 28 days each (day 0, day 28). ZYCOV-D® is approved for restricted use in emergency situation of COVID-19.

## **4.2 Posology and method of administration**

This vaccination schedule consists of 2 separate doses to be given at an interval of 28 days each (day 0 and day 28). Each dose consists of three shots of 0.1ml each given by needle free injector (Pharmajet Tropis device) via intradermal route at three separate sites (2 shots on one arm (recommended distance between two shots is at least 5 cms) and 1 shot on other arm).

### **Method of Administration:**

**ZYCOV-D<sup>®</sup>** has to be given by intradermal route only using needle free injector (Pharmajet Tropis device).

Kindly refer Medication Guide for step by step guidance on Method of Administration.

## **4.3 Contraindications**

**ZYCOV-D<sup>®</sup>** is contraindicated in individuals known to have hypersensitivity to the active substance or to any of the excipients.

## **4.4 Special warnings and precautions for use**

### ***Hypersensitivity***

As with all injectable vaccines, appropriate medical treatment and supervision should always be readily available in case of an anaphylactic event following the administration of the vaccine.

### ***Concurrent Illness***

As with other vaccines, administration of **ZYCOV-D<sup>®</sup>** should be postponed in individuals suffering from an acute severe febrile illness. However, the presence of a minor infection, such as cold, and/or low-grade fever should not delay vaccination.

### ***Immunocompromised individuals***

It is not known whether individuals with impaired immune responsiveness, including individuals receiving immunosuppressant therapy, will elicit the same response as immunocompetent individuals to the vaccine regimen. Immunocompromised individuals may have relatively weaker immune response to the vaccine regimen.

### ***Duration and level of protection***

The duration of protection has not yet been established. As with any vaccine, vaccination with ZYCOV-D<sup>®</sup> may not protect all vaccine recipients.

### ***Interchangeability***

No data are available on the use of ZYCOV-D<sup>®</sup> in persons that have previously received partial / complete vaccine series with another COVID-19 vaccine.

## **4.5 Interaction with other medicinal products and other forms of interaction**

No interaction studies have been performed. Concomitant administration of ZYCOV-D<sup>®</sup> with other vaccines has not been studied.

## **4.6 Special Population**

### ***Elderly Population:***

Efficacy and safety data are currently limited in individuals  $\geq 60$  years of age. No dosage adjustment is required in elderly individuals  $\geq 60$  years of age.

### ***Paediatric Population:***

Efficacy and safety data are currently limited in adolescents aged 12 to <18 years. The safety and efficacy of ZYCOV-D<sup>®</sup> in children (aged <12 years old) has not yet been established.

### ***Fertility***

There is no clinical data on the effect of ZYCOV-D<sup>®</sup> on fertility.

### ***Pregnancy***

The safety and efficacy of ZYCOV-D<sup>®</sup> in pregnancy has not been established.

### ***Breastfeeding***

The safety and efficacy of ZYCOV-D<sup>®</sup> in lactating females has not been established.

## **4.7 Effects on ability to drive and use machines**

ZYCOV-D<sup>®</sup> has no or negligible influence on the ability to drive and use machines. However, some of the adverse reactions may temporarily affect the ability to drive or use machines.

## 4.8 Undesirable effects

### Phase I/II Study:

A total of 1048 subjects were enrolled in the Phase I/II study, comprising of 4 different arms as follows:

- Arm 1: 1mg dose given by needle and syringe
- Arm 2: 1mg dose given by Pharmajet
- Arm 3: 2mg dose given by needle and syringe
- Arm 4: 2mg dose given by Pharmajet

**The age group and demographic characteristics of the subjects enrolled in Phase I/II study are as follows**

<b>Phase I: 48 adult subjects</b>				
	Arm 1, 1 mg (0.1 mL needle and syringe)	Arm 2, 1 mg (0.1 mL Pharmajet)	Arm 3, 2 mg (0.2 mL needle and syringe)	Arm 4, 2 mg (0.2 mL Pharmajet)
Adult subjects (Male)	12	12	12	12
Mean Age (Years)	35.4	31.8	35.1	37.2
<b>Phase II: 1000 subjects</b>				
	Arm 1, 1 mg (0.1 mL needle and syringe)	Arm 2, 1 mg (0.1 mL Pharmajet)	Arm 3, 2 mg (0.2 mL needle and syringe)	Arm 4, 2 mg (0.2 mL Pharmajet)
N	251	249	250	250
Male	188	186	179	177
Female	63	63	71	73
Adolescent	04	06	06	02
Mean Age (Years)	35.0 ± 11.83	34.2 ± 12.11	35.4 ± 10.46	34.6 ± 10.43

Adverse events reported in Phase I Study with 2mg Pharmajet Arm:

*Solicited adverse events:* Tenderness at the site of injection.

*Unsolicited adverse events:* Low WBC count.

Adverse events reported in Phase II Study with 2mg Pharmajet Arm:

Solicited adverse events: Nausea, fatigue, injection site erythema, injection site pain, injection site pruritus, injection site swelling, pyrexia, myalgia and headache.

**Frequency and Percentages of Participants with Solicited Local and systemic adverse events and unsolicited adverse events after each dose – Safety population**

AE Terms	ZyCoV-D n(%)			Placebo n(%)		
	Dose I (N = 200) n(%)	Dose II (N = 197) n(%)	Dose III (N = 194) n(%)	Dose I (N = 50) n(%)	Dose II (N = 49) n(%)	Dose III (N = 48) n(%)
<b>Solicited Local AEs</b>						
Pain at injection site	7 (3.50)	9 (4.57)	6 (3.09)	0 (0.00)	0 (0.00)	0 (0.00)
Redness at injection site	9 (4.50)	10 (5.08)	9 (4.64)	0 (0.00)	0 (0.00)	0 (0.00)
Swelling at injection site	5 (2.50)	6 (3.05)	5 (2.58)	0 (0.00)	0 (0.00)	0 (0.00)
Itching at injection site	1 (0.50)	7 (3.55)	2 (1.03)	0 (0.00)	0 (0.00)	0 (0.00)
Muscle pain	1 (0.50)	0 (0.00)	1 (0.52)	0 (0.00)	0 (0.00)	0 (0.00)
<b>Solicited Systemic AEs</b>						
Fatigue	3 (1.50)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	1 (2.08)
Fever	2 (1.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
Headache	3 (1.50)	1 (0.51)	0 (0.00)	1 (2.00)	0 (0.00)	1 (2.08)
Nausea	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	1 (2.08)
<b>Un Solicited Systemic AEs</b>						
Covid-19	2 (1.00)	0 (0.00)	2 (1.03)	0 (0.00)	0 (0.00)	1 (2.08)
Nasal Dryness	1 (0.50)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
High Blood Pressure	1 (0.50)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
Arthralgia	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
Body ache	0 (0.00)	3 (1.52)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
Chikungunya virus infection	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
Cough	0 (0.00)	2 (1.02)	1 (0.52)	0 (0.00)	0 (0.00)	0 (0.00)
Pyrexia	0 (0.00)	5 (2.54)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
Headache	0 (0.00)	2 (1.02)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
Myalgia	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
Rhinorrhoea	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
Asthenia	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
Dysuria	0 (0.00)	0 (0.00)	1 (0.52)	0 (0.00)	0 (0.00)	0 (0.00)
Fatigue	0 (0.00)	0 (0.00)	1 (0.52)	0 (0.00)	0 (0.00)	0 (0.00)
<b>N = number of subjects in the specified treatment arm; n = Number of participants with the specified event</b>						

Serious adverse events: 4 subjects experienced 7 serious adverse events: viral pneumonia, in-patient hospitalization [due to discharge at elbow site, pyrexia, arthralgia, joint swelling, erythema], surgical removal of orthopaedic implant, acute coronary syndrome, left ventricular failure with bronchopneumonia, and COVID-19 (02). None of these serious adverse events was related to IP.

All the adverse events reported in Phase I/II studies resolved without sequelae.

Adverse Events reported from Phase III Study - 2mg-3dose regimen (Interim data):

In our ongoing Phase III clinical trial, a total of 27703 subjects have been enrolled till the interim analysis. Amongst them more than 900 subjects belonged to the adolescent age group (12-17 years).

The age group and demographic characteristics of the subjects enrolled in Phase III study are as follows:

<b>Phase III: 27703 subjects (Data at the time of interim analysis), Total sample size: 28216</b>			
	Vaccine, 2 mg (0.2 mL Pharmajet)	Placebo	Total
Age (in Years) Mean	36.4	36.6	36.5
Age (12-17)	448	487	935
Age ( 18-60)	12364	12338	24702
Age (above 60)	1039	1027	2066
Gender			
Female	4506	4605	9111
Male	9345	9247	18592
Subjects at risk(Co-morbidities)	709	740	1449
Stable Chronic Heart Disease	167	155	322
Stable Chronic Lung Disease	13	7	20
Controlled Diabetic	275	289	564
Stable Liver Disease	2	3	5
Severe Obesity	18	14	32
Other Stable Co-morbid	295	293	588

The safety profile of the adolescent age group and the overall population has been found to be same. The common solicited and unsolicited adverse events reported in total population are as under:

*Solicited Local Adverse Events:* The most frequently reported solicited local adverse events across all treated subjects in both groups (ZYCOV-D<sup>®</sup> and Placebo) were pain at injection site: (0.66% and 0.62% after Dose 1; 0.34% and 0.35% after Dose 2 and 0.27% and 0.26% after Dose 3), redness: (0.31% and 0.28% after Dose 1; 0.19% and 0.09% after Dose 2 and 0.17% and 0.09% after Dose 3), swelling: (0.27% and 0.28% after Dose 1; 0.08% and 0.06% after Dose 2 and 0.09% and 0.05% after Dose 3) and itching: 0.08% and 0.14% after Dose 1; 0.05% and 0.07% after Dose 2 and 0.02% and 0.05% after Dose 3). Most of the adverse events were mild or moderate in severity. These events were comparable between ZYCOV-D<sup>®</sup> and placebo groups.

*Solicited Systemic Adverse Events:* The most commonly reported solicited systemic adverse events across all treated subjects in both groups (ZYCOV-D<sup>®</sup> and Placebo) were headache (0.25% and 0.22% after Dose 1; 0.20% and 0.24% after Dose 2 and 0.16% and 0.17% after Dose 3), fever (0.20% and 0.14% after Dose 1; 0.14% and 0.21% after Dose 2 and 0.13% and 0.10% after Dose 3), muscle pain (0.19% and 0.28% after Dose 1; 0.11% and 0.18% after Dose 2 and 0.11% and 0.09% after Dose 3), and fatigue (0.19% and 0.19% after Dose 1; 0.14% and 0.16% after Dose 2 and 0.09% and 0.13% after Dose 3). Most of the adverse events were mild or moderate in severity. These events were comparable between ZYCOV-D<sup>®</sup> and placebo groups.

Unsolicited Adverse Events: Arthralgia, Back pain, Muscle spasms, Myalgia, Musculoskeletal pain, Neck pain, Vertigo, Diarrhoea, Gastritis, Gastrooesophageal reflux disease, Nausea, Vomiting, Asthenia, Chills, Eye irritation, Abdominal distension, Abdominal pain, Fatigue, Pain, Pyrexia, Nasopharyngitis, Pain in extremity, Ageusia, Anosmia, Cerebral infarction, Dizziness, Headache, Cough, Dyspnoea, Nasal dryness, Oropharyngeal pain, Rhinorrhoea, Sneezing.

*Serious adverse events:* As per interim analysis report, 15 serious adverse events were identified: stroke (02), Death due to Cardiorespiratory arrest with septicaemia and alcoholic liver disease (1), Death due to COVID19 (1), Gram negative enteritis (1) and COVID19 (10). None of these serious adverse events was related to IP.

Adverse Events reported from Phase I/II Study - 3mg-2dose regimen:

A total of 150 adults healthy subjects >18 years of age were enrolled in the Phase I/II study of 3mg – 2 dose regimen (100 in Vaccine arm and 50 in the Placebo arm). The age group and demographic characteristics of the subjects enrolled in Phase I/II study are as follows:

	<b>Vaccine (N = 100)</b>	<b>Placebo (N = 50)</b>
<b>Mean Age (years)</b>	34.2	36.7
<b>Male</b>	68 (68.0%)	30 (60.0%)
<b>Female</b>	32 (32.0%)	20 (40.0%)

The following adverse events were reported during the study:

*Local site adverse events:* Pain (1.9%), Redness (1.4%), Swelling (1.2%), Itching (0.5%)

*Systemic adverse events:* Headache (2.6%), Tiredness / Fatigue (2.1%), Fever (1.1%), Diarrhea (0.5%) and Nausea (0.5%).

Adverse Events reported from Phase III Study - 3mg-2dose regimen:

In our ongoing Phase III clinical trial of 3mg – 2 dose regimens, a total of 3000 healthy subjects >12 years of age have been enrolled. Amongst them 1193 subjects belonged to the adolescent age group (12-17 years). The age group and demographic characteristics of the subjects enrolled in Phase III study are as follows:

	<b>Total (N = 3000)</b>	<b>Total (N = 1807)</b>	<b>Total (N = 1193)</b>
<b>Mean Age (Years)</b>	<b>27.4</b>	<b>35.9</b>	<b>14.6</b>
<b>Male</b>	<b>1907 (63.6%)</b>	<b>1278 (70.7%)</b>	<b>629 (52.7%)</b>
<b>Female</b>	<b>1093 (36.4%)</b>	<b>529 (29.3%)</b>	<b>564 (47.3%)</b>

The safety profile of the adolescent age group and the adult population has been found to be similar. The adverse events reported in total population up to day 112 are as under:

*Local site adverse events:* Pain (1.9%), Redness (0.7%), Swelling (0.4%), Itching (0.3%)

*Solicited systemic adverse events:* Headache (0.4%), Tiredness / Fatigue (0.2%), Fever (0.4%), arthralgia (0.2%), myalgia (0.1%) and vomiting (0.01%).

*Unsolicited systemic adverse events:* Body ache (0.13%), Weakness (0.12%), Headache (0.08%), Cough and Cold (0.08%), Fever (0.07%), Diarrhoea (0.05%), Tiredness (0.03%), COVID-19 (0.06%), Myalgia (0.01%) and Vomiting (0.01%).

## 4.9 Overdose

Experience of overdose is limited.

There is no specific treatment for an overdose with ZYCOV-D®. In the event of an overdose, the individual should be monitored and provided with symptomatic treatment as appropriate.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Mechanism of Action

The plasmid construct of ZYCOV-D® carrying the spike-S gene of interest enters host cells, where it remains in the nucleus as an episome; without getting integrated into the host cell DNA. Thus, using the host cell's protein translation machinery, the inserted cloned gene in the episome will direct the synthesis of the antigen it encodes. The protein produced by plasmid-transfected cells is likely to be expressed within the cell and folded in its native conformation. Further the signal peptide prompts cells to translocate the protein, usually to the cellular membrane. The antigen is recognized by antigen presenting cells (APCs) and further induces antibodies including neutralizing antibodies and cellular immune response through major histocompatibility complex (MHC).

### 5.2 Pharmacodynamics properties

Immunogenicity Data 28 days after last dose from Phase II and Phase III Clinical Trials (2mg-3dose regimen with Pharmajet):

#### Phase II Clinical Trial:

Parameter	Data
Seroconversion rate based on IgG* (%)	91.28%
Seroconversion rate based on Neutralizing Antibody response^ (%)	88.89%
GMT based on Neutralizing Antibody response^	131.32 (63.50, 271.58) <sup>§</sup>
GMFR based on Neutralizing Antibody response^	22.56 (10.57, 48.16) <sup>§</sup>
*by S1 antigen ELISA	
^Wild type virus neutralization assay (PRNT <sub>50</sub> )	
§ data presented as Geometric Mean (95% CI)	

### Phase III Clinical Trial:

Parameter	Data
Seroconversion rate based on IgG* (%)	93.33%
GMT based on IgG*	952.67 (707.9, 1282.0) <sup>\$</sup>
GMFR based on IgG*	136.09 (101.11, 183.1) <sup>\$</sup>
*by S1 antigen ELISA \$ data presented as Geometric Mean (95% CI)	

#### Interim Efficacy Data from Phase III Clinical Trial (2mg-3dose regimen with Pharmajet):

A total of 27703 subjects were enrolled in the Phase III study till interim analysis. The interim primary efficacy analysis was based on the Per-Protocol analysis, which consisted of all participants with negative baseline SARS-CoV-2 status (i.e., negative RT-PCR for SARS-CoV-2) and who had received 3 doses of investigational product. Total of 12350 subjects who had completed 84±3 days in vaccine group and total 12320 subjects who had completed 84±3 days in placebo group were considered for analysis. Out of 81 symptomatic RT-PCR positive COVID-19 cases considered for interim analysis, 61 were in placebo group and 20 were in the vaccine (ZYCOV-D<sup>®</sup>) group. On the basis of calculation, ZYCOV-D<sup>®</sup> vaccine efficacy is 66.6% (95% CI: 47.6 to 80.7).

#### Immunogenicity Data 28 days after second dose from Phase III Clinical Trial (3mg-2dose regimen with Pharmajet) in subjects seronegative at baseline:

Parameter	Total	Adults	Adolescents
Seroconversion rate based on IgG* (%)	95.3%	96.3%	96.3%
GMT based on IgG*	1262.9 (960.0 to 1661.4)	1088.2 (736.2 to 1608.5)	1465.7 (990.3 to 2169.3)
GMFR based on IgG*	180.4 (137.2 to 237.4)	155.5 (105.2 to 229.8)	209.4 (141.5 to 309.9)
*by S1 antigen ELISA \$ data presented as Geometric Mean (95% CI)			

### 5.3 Pharmacokinetic properties

- Not applicable

## **6. Preclinical safety data**

### **6.1 Animal Pharmacology:**

The immunogenicity potential of ZYCOV-D<sup>®</sup> has been evaluated in mice, guinea pig and rabbit models by intradermal route at varying dose levels. Immunogenicity studies in animals demonstrated that the candidate DNA vaccine induces robust antibody response including neutralizing antibodies against SARS-CoV-2 and also provided Th-1 response as evidenced by elevated IFN- $\gamma$  levels. In animal studies primary antibody response starts mounting in serum two weeks after two doses and reaches peak two weeks after third immunization. The serum IgG levels against spike antigen in mice were maintained even after three months post last dosing suggesting a long-term immune response generated by the DNA vaccine candidate.

Protective efficacy of ZYCOV-D<sup>®</sup> was also evaluated in Rhesus Macaques. We assessed the immunogenicity and protective efficacy of two formulations (1mg and 2mg) of ZYCOV-D<sup>®</sup> administered either through Needle Free Injection System (NFIS) and syringe needle (intradermal) with three dose vaccine regimens. ZYCOV-D<sup>®</sup> demonstrated good immunogenicity as can be seen by the analysis of SARS-CoV-2 specific IgG (S1), Neutralizing Antibody (Nab) titres, percentage lymphocytes and cytokines response during immunization and after virus challenge. The viral clearance in nasal swab (NS), throat swab (TS), and bronchoalveolar lavage (BAL) in animals receiving ZYCOV-D<sup>®</sup> was seen demonstrating protective efficacy.

### **6.2 Animal Toxicology**

Non-clinical data reveal no special hazard for humans based on a conventional study of repeat dose toxicity. Animal studies evaluating potential toxicity to reproduction and development have not yet been completed.

28-day repeat dose preclinical toxicology (PCT) studies were conducted in Wistar rats and New Zealand white rabbits and the vaccine was found to be safe and well-tolerated. Indeed, no treatment related adverse effects and behavioral changes were observed in animals during the studies. Further, histopathological examination reveals no changes of toxicological significance at high dose of 3mg (1.5 times the intended single human dose) and 6mg (3 times the intended single human dose) in rats and rabbits respectively.

## **7. Description**

**ZYCOV-D<sup>®</sup>** is a DNA based vaccine for prevention of COVID-19. It comprises of a DNA plasmid vector carrying full length spike (S) gene region expressing SARS-CoV-2 spike (S) protein along with gene coding for signal peptide. The spike gene region was selected from submitted Wuhan Hu-1 isolate sequence (Genebank Accession No. MN908947.3). The S protein of the virus includes the receptor binding domain (RBD), responsible for binding to the human angiotensin converting enzyme-2 (ACE-2) receptor, which mediates the entry of virus inside the cell. The DNA plasmid construct was transformed into E. coli cells for large scale production.

## **8. PHARMACEUTICAL PARTICULARS**

### **8.1 List of excipients**

Not Applicable, as no excipient is being used.

### **8.2 Incompatibilities**

This vaccine should not be mixed with any other medicinal product.

### **8.3 Shelf life**

The expiry date of vaccine is indicated on the label and packaging. Once opened, multi-dose vials should be used as soon as practically possible and within 6 hours when kept between +2°C and +8°C. All opened multidose vials of ZYCOV-D<sup>®</sup> should be discarded at the end of immunization session or within 6 hours whichever comes first.

### **8.4 Special precautions for storage**

Store at 2° to 8°C. Do Not Freeze. In case of unexpected freezing of vaccine at 2-8°C storage, it can be administered after thawing.

Multidose Vials: To be used within 6 hours of opening.

### **8.5 Packing information**

ZYCOV-D<sup>®</sup> is supplied in a USP type-1 tubular glass vial 2.0 ml.

## **8.6 Special precautions for disposal**

Any unused product or waste material should be disposed of in accordance with local requirements.

## **9. Details of manufacturer**

### **Zydus Lifesciences Limited**

(formerly known as Cadila Healthcare Limited)

Plot Survey No. 23, 25/P, 37, 40/P, 42 to 47

Sarkhej- Bavla N.H. 8A, Opp. Ramdev Masala,

Village: Changodar, Taluka: Sanand,

Dist. Ahmedabad – 382 213

Phone: +91-2717- 664600

## **10. MARKETING AUTHORISATION NUMBER(S)**

MF/BIO/22/000034

## **11. DATE OF FIRST AUTHORISATION**

25-Apr-2022



**ARTWORK APPROVAL FORM**

<b>PRODUCT NAME:</b>	Zycov-D		<b>ITEM CODE:</b>	2083605	
			<b>SUPERSEDED ITEM CODE:</b>	NA	
			<b>SPECIFICATION NO:</b>	QC/SPC/P/7045	
<b>APPLICABLE FINISHED GOODS:</b>	5040666		<b>COUNTRY:</b>	INDIA	
<b>COMPONENT:</b>	LITERATURE	<b>PRIMARY PACK:</b>	20 X 2 ml (Multi Dose) Vial	<b>MARKET:</b>	India
<b>ACTUAL SIZE:</b>	128 x 220 mm (LxH)	<b>ARTWORK SIZE:</b>	100%		
<b>SUBSTRATE:</b>	60 GSM Maplitho Paper	<b>FORM OF SUPPLY:</b>	Folded Form : 64X55 mm ± 2 mm ( L x H )		
<b>PHARMACODE VALUE:</b>	17	<b>BARCODE NO:</b>	NA		
<b>COLOR SCHEME:</b>	One Colour	<b>PANTONE SHADES:</b>	Black		
<b>CHANGE CONTROL NUMBER:</b>	NA	<b>REASON FOR CHANGES:</b>	New		
<b>SPECIAL INSTRUCTIONS (IF ANY):</b>	1. Mfg. Site: Vaccine Technology Centre				

Date.:14.06.22

\\sigma-file-clu\ptc packaging\Design Studio\01 Artworks Store\Domestic\01 Zydus Lifesciences\ZLL\Zycov-D Vaccine\2083605\_LIT. Zycov-D Vaccine 2ml Vial SL DOM\2083605

Page : 2 of 2

0 and day 28). Each dose consists of three shots of 0.1ml each given by needle free injector (Pharmajet Tropis device) via intradermal route at three separate sites (2 shots on one arm (recommended distance between two shots is at least 5 cms) and 1 shot on other arm).

**ZYCOV-D<sup>®</sup>** vaccine has shown efficacy of 66.6% in interim analysis of Phase III clinical trial with 2 mg - 3dose regimen. This vaccine may not protect everyone.

**WHAT YOU NEED TO KNOW BEFORE YOU GET THIS VACCINE**

**WHAT IS COVID-19?**

COVID-19 disease is caused by a coronavirus called SARS-CoV-2. This type of coronavirus has not been seen before. You can get COVID-19 through contact with another person who has the virus. It is predominantly a respiratory illness that can affect other organs. People with COVID-19 have had a wide range of symptoms reported, ranging from mild symptoms to severe illness. Symptoms may appear 2 to 14 days after exposure to the virus. Symptoms may include: fever or chills; cough; shortness of breath; fatigue; muscle or body aches; headache; new loss of taste or smell; sore throat; congestion or runny nose; nausea or vomiting; diarrhea.

**WHAT IS ZYCOV-D<sup>®</sup> VACCINE?**

**ZYCOV-D<sup>®</sup>** is approved for restricted use in emergency situation vaccine that may prevent COVID-19 disease in individuals 12 years of age and older.

**WHAT SHOULD YOU MENTION TO YOUR HEALTHCARE PROVIDER BEFORE YOU GET ZYCOV-D<sup>®</sup> VACCINE?**

**Tell the healthcare provider about all of your medical conditions, including:**

- If you have ever had a severe allergic reaction (anaphylaxis) after any drug, food, any vaccine or any ingredients of **ZYCOV-D<sup>®</sup>** vaccine
- If you have fever
- If you have a bleeding disorder or are on a blood thinner
- If you are immunocompromised or are on a medicine that affects your immune system
- If you are pregnant or plan to become pregnant
- If you are breastfeeding
- If you have received another COVID-19 vaccine

You should consult your healthcare provider before deciding to take the vaccine.

**WHO SHOULD GET ZYCOV-D<sup>®</sup> VACCINE?**

**ZYCOV-D<sup>®</sup>** Vaccine has been approved for restricted use in emergency situation in individuals 12 years of age and older.

**WHO SHOULD NOT GET ZYCOV-D<sup>®</sup> VACCINE?**

- You should not get **ZYCOV-D<sup>®</sup>** Vaccine if you:
- had a severe allergic reaction after a previous dose of this vaccine
  - had a severe allergic reaction to any ingredient of this vaccine
  - had a severe allergic reaction to any other vaccine

**WHAT ARE THE INGREDIENTS IN ZYCOV-D<sup>®</sup> VACCINE?**

**ZYCOV-D<sup>®</sup>** Vaccine includes the following ingredients: Deoxyribonucleic acid (DNA)  
Phosphate buffered saline

**HOW IS ZYCOV-D<sup>®</sup> GIVEN?**

**ZYCOV-D<sup>®</sup>** Vaccine will be given to you as an Intradermal (ID) injection only using Needle Free Injector (Pharmajet Tropis device).

**ZYCOV-D<sup>®</sup>** vaccination course consists of 2 separate doses of 3mg each.

If you receive one dose of **ZYCOV-D<sup>®</sup>** vaccine, then the 2<sup>nd</sup> dose should be administered 28 days after the previous dose.

**If you miss your 2<sup>nd</sup> dose**

If you forget to go back at the scheduled time, ask your healthcare provider for advice. It is important that you return for your 2<sup>nd</sup> dose of **ZYCOV-D<sup>®</sup>** vaccine as per your dosing regimen.

**HAS ZYCOV-D<sup>®</sup> VACCINE BEEN USED BEFORE?**

**ZYCOV-D<sup>®</sup>** is being used in clinical trials, a number of participants have received one or two or three doses in trials being conducted in India.

**WHAT ARE THE BENEFITS OF ZYCOV-D<sup>®</sup> VACCINE?**

In ongoing clinical trials, **ZYCOV-D<sup>®</sup>** Vaccine has been shown to prevent COVID-19 disease following 3 doses of 2mg each given 28 days apart. The duration of protection against COVID-19 disease is currently unknown.

You may get protective immune response 28 days after the 3<sup>rd</sup> dose of 2mg - 3 dose regimen of **ZYCOV-D<sup>®</sup>** vaccine. Similar immune response is also seen 28 days after 2nd dose of 3mg - 2 dose regimen of ZyCoV-D vaccine. The comparison of immune response of 2mg - 3 dose regimen and 3mg - 2 dose regimen is shown in table below:

Table: Immunogenicity Data in subjects seronegative at baseline:

Statistics	Cohort	Day 56 Data of 3mg - 2 dose regimen	Day 84 data of 2mg - 3 dose regimen
Seroconversion rate based on IgG* (%)	<b>Total</b>	122 / 128 (95.3%)	168 / 180 (93.3%)
	<b>Adults</b>	61/64 (95.3%)	132/142 (93.0%)
	<b>Pediatric</b>	61/64 (95.3%)	36/38 (94.7%)
GMT based on IgG*	<b>Total</b>	1262.9 (960.0 to 1661.4)	998.1 (778.7 to 1279.3)
	<b>Adults</b>	1088.2 (736.2 to 1608.5)	926.0 (693.9 to 1235.9)
	<b>Pediatric</b>	1465.7 (990.3 to 2169.3)	1320.9 (816.4 to 2137.1)

\*by S1 antigen ELISA

\$ data presented as Geometric Mean (95% CI)

**WHAT ARE THE RISKS OF ZYCOV-D<sup>®</sup> VACCINE?**

Common side effects that have been reported with **ZYCOV-D<sup>®</sup>** Vaccine include:

- Injection site redness
- Injection site pain
- Injection site itching
- Injection site swelling
- Fever
- Muscle Pain
- Headache
- Nausea
- Fatigue / Tiredness
- Diarrhea

These may not be all the possible side effects of **ZYCOV-D<sup>®</sup>** Vaccine. Serious and unexpected side effects may occur. **ZYCOV-D<sup>®</sup>** Vaccine is still being studied in clinical trials.

**WHAT SHOULD I DO ABOUT SIDE EFFECTS?**

If you experience a severe allergic reaction, call or go to the nearest hospital.

Call the healthcare provider if you have any side effects that bother you or do not go away.

In addition, you can report side effects after vaccination to Zydus Lifesciences Limited who is the manufacturer of **ZYCOV-D<sup>®</sup>** vaccine on toll free number 1800 419 1141 or visit www.zyduslife.com

## ARTWORK APPROVAL FORM

<b>PRODUCT NAME:</b>	Zycov-D	<b>ITEM CODE:</b>	2083603
		<b>SUPERSEDED ITEM CODE:</b>	NA
		<b>SPECIFICATION NO:</b>	QC/SPC/P/7045
<b>APPLICABLE FINISHED GOODS:</b>	5040666	<b>COUNTRY:</b>	INDIA
<b>COMPONENT:</b>	PACKINSERT	<b>PRIMARY PACK:</b>	20 X 2 ml (Multi Dose) Vial
<b>ACTUAL SIZE:</b>	128 x 232 mm (L x H)	<b>ARTWORK SIZE:</b>	100%
<b>SUBSTRATE:</b>	60 GSM Maplitho Paper	<b>FORM OF SUPPLY:</b>	Folded Form : 64X58 mm ± 2 mm ( L x H )
<b>PHARMACODE VALUE:</b>	56	<b>BARCODE NO:</b>	NA
<b>COLOR SCHEME:</b>	P 2685 C	<b>PANTONE SHADES:</b>	1 Colour
<b>CHANGE CONTROL NUMBER:</b>	NA	<b>REASON FOR CHANGES:</b>	New
<b>SPECIAL INSTRUCTIONS (IF ANY):</b>	Mfg. Site: Vaccine Technology Centre		

Date.:14.06.22

\\sigma-file-clu\ptc packaging\Design Studio\01 Artworks Store\Domestic\01 Zydus Lifesciences\ZLL\Zycov-D Vaccine\2083603\_PI Zycov-D Vaccine 2ml Vial SL DOM\2083603

Page : 1 of 2

group and 20 were in the vaccine (ZYCOV-D<sup>®</sup>) group. On the basis of calculation, ZYCOV-D<sup>®</sup> vaccine efficacy is 66.6% (95% CI: 47.6 to 80.7). *Immunogenicity Data 28 days after second dose from Phase III Clinical Trial (3mg-2dose regimen with Pharmajet) in subjects seronegative at baseline:*

Parameter	Total	Adults	Adolescents
Seroconversion rate based on IgG* (%)	95.3%	95.3%	95.3%
GMT based on IgG*	1262.9 (960.0 to 1661.4)	1088.2 (736.2 to 1608.5)	1465.7 (990.3 to 2169.3)
GMFR based on IgG*	180.4 (137.2 to 237.4)	155.5 (105.2 to 229.8)	209.4 (141.5 to 309.9)

\*by S1 antigen ELISA  
§ data presented as Geometric Mean (95% CI)

### 5.3 Pharmacokinetic properties

NA

### 6. Nonclinical properties

#### 6.1 Animal Pharmacology

The immunogenicity potential of ZYCOV-D<sup>®</sup> has been evaluated in mice, guinea pig and rabbit models by intradermal route at varying dose levels. Immunogenicity studies in animals demonstrated that the candidate DNA vaccine induces robust antibody response including neutralizing antibodies against SARS-CoV-2 and also provided Th-1 response as evidenced by elevated IFN-γ levels. In animal studies primary antibody response starts mounting in serum two weeks after two doses and reaches peak two weeks after third immunization. The serum IgG levels against spike antigen in mice were maintained even after three months post last dosing suggesting a long-term immune response generated by the DNA vaccine candidate. Protective efficacy of ZYCOV-D<sup>®</sup> was also evaluated in Rhesus Macaques. We assessed the immunogenicity and protective efficacy of two formulations (1mg and 2mg) of ZYCOV-D<sup>®</sup> administered either through Needle Free Injection System (NFIS) and syringe needle (intradermal) with three dose vaccine regimens. ZYCOV-D<sup>®</sup> demonstrated good immunogenicity as can be seen by the analysis of SARS-CoV-2 specific IgG (S1), Neutralizing Antibody (NaB) titres, percentage lymphocytes and cytokines response during immunization and after virus challenge. The viral clearance in nasal swab (NS), throat swab (TS), and bronchoalveolar lavage (BAL) in animals receiving ZYCOV-D<sup>®</sup> was seen demonstrating protective efficacy.

#### 6.2 Animal Toxicology

Non-clinical data reveal no special hazard for humans based on a conventional study of repeat dose toxicity. Animal studies evaluating potential toxicity to reproduction and development have not yet been completed.

28-day repeat dose preclinical toxicology (PCT) studies were conducted in Wistar rats and New Zealand white rabbits and the vaccine was found to be safe and well-tolerated. Indeed, no treatment related adverse effects and behavioural changes were observed in animals during the studies. Further, histopathological examination reveals no changes of toxicological significance at high dose of 3mg (1.5 times the intended single human dose) and 6mg (3 times the intended single human dose) in rats and rabbits respectively.

### 7. Description

ZYCOV-D<sup>®</sup> is a DNA based vaccine for prevention of COVID-19. It comprises of a DNA plasmid vector carrying full length spike (S) gene region expressing SARS-CoV-2 spike (S) protein along with gene coding for signal peptide. The spike gene region was selected from submitted Wuhan Hu-1 isolate sequence (Genebank Accession No. MN908947.3). The S protein of the virus includes the receptor binding domain (RBD), responsible for binding to the human angiotensin converting enzyme-2 (ACE-2) receptor, which mediates the entry of virus inside the cell. The DNA plasmid construct was transformed into E. coli cells for large scale production.

### 8. Pharmaceutical particulars

#### 8.1 Incompatibilities

This vaccine should not be mixed with any other medicinal product.

#### 8.2 Shelf-life

The expiry date of vaccine is indicated on the label and packaging. Once opened, multi-dose vials should be used as soon as practically possible and within 6 hours when kept between +2°C and +8°C. All opened multidose vials of ZYCOV-D<sup>®</sup> should be discarded at the end of immunization session or within 6 hours whichever comes first.

#### 8.3 Packaging information

ZYCOV-D<sup>®</sup> is supplied in a USP type-1 tubular glass vial 2.0 mL.

#### 8.4 Storage and handling instructions

Store at 2° to 8°C. Do Not Freeze. In case of unexpected freezing of vaccine at 2-8°C storage, it can be administered after thawing. Multidose Vials: To be used within 6 hours of opening

### 9. Details of manufacturer

**Zydus Lifesciences Limited**  
(formerly known as Cadila Healthcare Limited)  
Plot Survey No.: 23, 25/P, 37, 40/P, 42 to 47,  
Sarkhej-Bavla N.H. No. 8A, Changodar, Tal: Sanand,  
Dist.: Ahmedabad-382213, Gujarat

### 10. Details of permission or licence number with date

MF/BIO/22/000034 Dated 25-Apr-2022

### 11. Date of revision

13/06/2022



To report adverse events, call toll free on  
1800 419 1141 or visit www.zyduslife.com  
© Registered Trademark



Approved for restricted use in emergency situation of COVID-19. For the use of a Registered Medical Practitioner or a Hospital or a laboratory only.

### CAUTION

3mg – 2 Dose Regimen: 3 shots of 0.1mL each should be given on day 0 and 28  
3x0.1 mL 3x0.1 mL  
Day 0 Day 28

It is recommended to use the vaccine only with Pharmajet Device. Using it with conventional needle and syringe will not lead to optimal immunogenicity response and will affect the efficacy of the vaccine.

### 1. Name of Medicinal Product

Novel Corona Virus 2019 - nCoV Vaccine  
(Recombinant)

### ZYCOV-D<sup>®</sup>

### 2. Qualitative and quantitative composition

Each 0.1 mL contains:  
DNA plasmid construct with spike protein gene region  
from SARS-CoV-2 virus Produced in E.coli 1.0 mg  
Phosphate Buffered saline q.s.

### 3. Dosage form and strength

Solution for Intradermal Injection.

Each dose consists of three shots of 0.1 mL each

### 4. Clinical particulars

#### 4.1 Therapeutic indication

ZYCOV-D<sup>®</sup> is indicated for active immunisation to prevent COVID-19 caused by SARS-CoV-2 in individuals 12 years of age and older when given in two separate doses of 3mg (0.3ml) each to be given at an interval of 28 days each (day 0, day 28). ZYCOV-D<sup>®</sup> is approved for restricted use in emergency situation of COVID-19.

#### 4.2 Posology and method of administration

This vaccination schedule consists of 2 separate doses to be given at an interval of 28 days each (day 0 and day 28). Each 3mg dose consists of three shots of 0.1mL each given by needle free injector (Pharmajet Tropis device) via intradermal route at three separate sites (2 shots on one arm (recommended distance between two shots is at least 5 cms) and 1 shot on other arm).

#### Method of Administration:

ZYCOV-D<sup>®</sup> has to be given by intradermal route only using needle free injector (Pharmajet Tropis device).

#### Kindly refer Medication Guide for step by step guidance on Method of Administration

### 4.3 Contraindications

ZYCOV-D<sup>®</sup> is contraindicated in individuals known to have hypersensitivity to the active substance or to any of the excipients

### 4.4 Special warnings and precautions for use

#### Hypersensitivity

As with all injectable vaccines, appropriate medical treatment and supervision should always be readily available in case of an anaphylactic event following the administration of the vaccine.

#### Concurrent illness

As with other vaccines, administration of ZYCOV-D<sup>®</sup> should be postponed in individuals suffering from an acute severe febrile illness. However, the presence of a minor infection, such as cold, and/or low grade fever should not delay vaccination.

#### Immunocompromised individuals

It is not known whether individuals with impaired immune responsiveness, including individuals receiving immunosuppressant therapy, will elicit the same response as immunocompetent individuals to the vaccine regimen. Immunocompromised individuals may have relatively weaker immune response to the vaccine regimen.

#### Duration and level of protection

The duration of protection has not yet been established. As with any vaccine, vaccination with ZYCOV-D<sup>®</sup> may not protect all vaccine recipients.

#### Interchangeability

No data are available on the use of ZYCOV-D<sup>®</sup> in persons that have previously received partial / complete vaccine series with another COVID-19 vaccine.

### 4.5 Interactions

No interaction studies have been performed. Concomitant administration of ZYCOV-D<sup>®</sup> with other vaccines has not been studied

### 4.6 Use in special populations (such as pregnant women, lactating women, paediatric patients, geriatric patients etc.)

#### Elderly Population:

Efficacy and safety data are currently limited in individuals ≥ 60 years of age. No dosage adjustment is required in elderly individuals ≥ 60 years of age.

#### Paediatric Population:

Efficacy and safety data are currently limited in adolescents aged 12 to <18 years. The safety and efficacy of ZYCOV-D<sup>®</sup> in children (aged <12 years old) has not yet been established.

#### Fertility

There is no clinical data on the effect of ZYCOV-D<sup>®</sup> on fertility.

#### Pregnancy

The safety and efficacy of ZYCOV-D<sup>®</sup> in pregnancy has not been established.

#### Breastfeeding

The safety and efficacy of ZYCOV-D<sup>®</sup> in lactating females has not been established.

### 4.7 Effects on ability to drive and use machines

ZYCOV-D<sup>®</sup> has no or negligible influence on the ability to drive and use machines. However, some of the adverse reactions may temporarily affect the ability to drive or use machines.

### 4.8 Undesirable effects

#### Phase I/II Study:

A total of 1048 subjects were enrolled in the Phase I/II study, comprising of 4 different arms as follows:

- Arm 1: 1mg dose given by needle and syringe
- Arm 2: 1mg dose given by Pharmajet
- Arm 3: 2mg dose given by needle and syringe
- Arm 4: 2mg dose given by Pharmajet

*The age group and demographic characteristics of the subjects enrolled*

**ARTWORK APPROVAL FORM**

<b>PRODUCT NAME:</b>	Zycov-D	<b>ITEM CODE:</b>	2083603	
		<b>SUPERSEDED ITEM CODE:</b>	NA	
		<b>SPECIFICATION NO:</b>	QC/SPC/P/7045	
<b>APPLICABLE FINISHED GOODS:</b>	5040666	<b>COUNTRY:</b>	INDIA	
<b>COMPONENT:</b>	PACKINSERT	<b>PRIMARY PACK:</b>	20 X 2 ml (Multi Dose) Vial	<b>MARKET:</b> India
<b>ACTUAL SIZE:</b>	128 x 232 mm (L x H)	<b>ARTWORK SIZE:</b>	100%	
<b>SUBSTRATE:</b>	60 GSM Maplitho Paper	<b>FORM OF SUPPLY:</b>	Folded Form : 64X58 mm ± 2 mm ( L x H )	
<b>PHARMACODE VALUE:</b>	56	<b>BARCODE NO:</b>	NA	
<b>COLOR SCHEME:</b>	P 2685 C	<b>PANTONE SHADES:</b>	1 Colour	
<b>CHANGE CONTROL NUMBER:</b>	NA	<b>REASON FOR CHANGES:</b>	New	
<b>SPECIAL INSTRUCTIONS (IF ANY):</b>	Mfg. Site: Vaccine Technology Centre			

Date: 14.06.22

\\sigma-file-clu\ptc packaging\Design Studio\01 Artworks Store\Domestic\01 Zydus Lifesciences\ZLL\Zycov-D Vaccine\2083603\_PI Zycov-D Vaccine 2ml Vial SL DOM\2083603

Page : 2 of 2

<p><i>in Phase I/II study are as follows:</i></p> <table border="1"> <thead> <tr> <th colspan="5">Phase I: 48 adult subjects</th> </tr> <tr> <th></th> <th>Arm 1, 1 mg (0.1 mL needle and syringe)</th> <th>Arm 2, 1 mg (0.1 mL Pharmajet)</th> <th>Arm 3, 2 mg (0.2 mL needle and syringe)</th> <th>Arm 4, 2 mg (0.2 mL Pharmajet)</th> </tr> </thead> <tbody> <tr> <td>Adult subjects (Male)</td> <td>12</td> <td>12</td> <td>12</td> <td>12</td> </tr> <tr> <td>Mean Age (Years)</td> <td>35.4</td> <td>31.8</td> <td>35.1</td> <td>37.2</td> </tr> </tbody> </table> <table border="1"> <thead> <tr> <th colspan="5">Phase II: 1000 subjects</th> </tr> <tr> <th></th> <th>Arm 1, 1 mg (0.1 mL needle and syringe)</th> <th>Arm 2, 1 mg (0.1 mL Pharmajet)</th> <th>Arm 3, 2 mg (0.2 mL needle and syringe)</th> <th>Arm 4, 2 mg (0.2 mL Pharmajet)</th> </tr> </thead> <tbody> <tr> <td>N</td> <td>251</td> <td>249</td> <td>250</td> <td>250</td> </tr> <tr> <td>Male</td> <td>188</td> <td>186</td> <td>179</td> <td>177</td> </tr> <tr> <td>Female</td> <td>63</td> <td>63</td> <td>71</td> <td>73</td> </tr> <tr> <td>Adolescent</td> <td>04</td> <td>06</td> <td>06</td> <td>02</td> </tr> <tr> <td>Mean Age (Years)</td> <td>35.0 ± 11.83</td> <td>34.2 ± 12.11</td> <td>35.4 ± 10.46</td> <td>34.6 ± 10.43</td> </tr> </tbody> </table> <p><b>Adverse events reported in Phase I Study with 2mg Pharmajet Arm:</b> Solicited adverse events: Tenderness at the site of injection. Unsolicited adverse events: Low WBC count.</p> <p><b>Adverse events reported in Phase II Study with 2mg Pharmajet Arm:</b> Solicited adverse events: Nausea, fatigue, injection site erythema, injection site pain, injection site pruritus, injection site swelling, pyrexia, myalgia and headache.</p> <p><b>Frequency and Percentages of Participants with Solicited Local and systemic adverse events and Unsolicited adverse events after each dose – Safety population</b></p> <table border="1"> <thead> <tr> <th rowspan="2">AE Terms</th> <th colspan="3">ZyCoV-D n(%)</th> <th colspan="3">Placebo n(%)</th> </tr> <tr> <th>Dose I (N = 200) n(%)</th> <th>Dose II (N = 197) n(%)</th> <th>Dose III (N = 194) n(%)</th> <th>Dose I (N = 50) n(%)</th> <th>Dose II (N = 49) n(%)</th> <th>Dose III (N = 48) n(%)</th> </tr> </thead> <tbody> <tr> <td colspan="7"><b>Solicited Local AEs</b></td> </tr> <tr> <td>Pain at injection site</td> <td>7 (3.50)</td> <td>9 (4.57)</td> <td>6 (3.09)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Redness at injection site</td> <td>9 (4.50)</td> <td>10 (5.08)</td> <td>9 (4.64)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Swelling at injection site</td> <td>5 (2.50)</td> <td>6 (3.05)</td> <td>5 (2.58)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Itching at injection site</td> <td>1 (0.50)</td> <td>7 (3.55)</td> <td>2 (1.03)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Muscle pain</td> <td>1 (0.50)</td> <td>0 (0.00)</td> <td>1 (0.52)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td colspan="7"><b>Solicited Systemic AEs</b></td> </tr> <tr> <td>Fatigue</td> <td>3 (1.50)</td> <td>1 (0.51)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>1 (2.08)</td> </tr> <tr> <td>Fever</td> <td>2 (1.00)</td> <td>1 (0.51)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Headache</td> <td>3 (1.50)</td> <td>1 (0.51)</td> <td>0 (0.00)</td> <td>1 (2.00)</td> <td>0 (0.00)</td> <td>1 (2.08)</td> </tr> <tr> <td>Nausea</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>1 (2.08)</td> </tr> <tr> <td colspan="7"><b>Un Solicited Systemic AEs</b></td> </tr> <tr> <td>Covid-19</td> <td>2 (1.00)</td> <td>0 (0.00)</td> <td>2 (1.03)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>1 (2.08)</td> </tr> <tr> <td>Nasal Dryness</td> <td>1 (0.50)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>High Blood Pressure</td> <td>1 (0.50)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Arthralgia</td> <td>0 (0.00)</td> <td>1 (0.51)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Body ache</td> <td>0 (0.00)</td> <td>3 (1.52)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Chikungunya virus infection</td> <td>0 (0.00)</td> <td>1 (0.51)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Cough</td> <td>0 (0.00)</td> <td>2 (1.02)</td> <td>1 (0.52)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Pyrexia</td> <td>0 (0.00)</td> <td>5 (2.54)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Headache</td> <td>0 (0.00)</td> <td>2 (1.02)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Myalgia</td> <td>0 (0.00)</td> <td>1 (0.51)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Rhinorrhoea</td> <td>0 (0.00)</td> <td>1 (0.51)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Asthenia</td> <td>0 (0.00)</td> <td>1 (0.51)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Dysuria</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>1 (0.52)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> <tr> <td>Fatigue</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>1 (0.52)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> <td>0 (0.00)</td> </tr> </tbody> </table> <p>N = number of subjects in the specified treatment arm; n = Number of participants with the specified event</p> <p><b>Serious adverse events:</b> 4 subjects experienced 7 serious adverse events: viral pneumonia, in-patient hospitalization [due to discharge at elbow site], pyrexia, arthralgia, joint swelling, erythema, surgical removal of orthopaedic implant, acute coronary syndrome, left ventricular failure with bronchopneumonia, and COVID-19 (02). None of these serious adverse events was related to IP.</p> <p>All the adverse events reported in Phase I/II studies resolved without sequelae.</p> <p><b>Adverse Events reported from Phase III Study - 2mg-3dose regimen (Interim data):</b> In our ongoing Phase III clinical trial, a total of 27703 subjects have been enrolled till the interim analysis. Amongst them more than 900 subjects belonged to the adolescent age group (12-17 years). The age group and demographic characteristics of the subjects enrolled in Phase III study are as follows:</p> <table border="1"> <thead> <tr> <th colspan="4">Phase III: 27703 subjects (Data at the time of interim analysis), Total sample size: 28216</th> </tr> <tr> <th></th> <th>Vaccine, 2 mg (0.2 mL Pharmajet)</th> <th>Placebo</th> <th>Total</th> </tr> </thead> <tbody> <tr> <td>Age (in Years) Mean</td> <td>36.4</td> <td>36.6</td> <td>36.5</td> </tr> <tr> <td>Age (12-17)</td> <td>448</td> <td>487</td> <td>935</td> </tr> <tr> <td>Age (18-60)</td> <td>12364</td> <td>12338</td> <td>24702</td> </tr> <tr> <td>Age (above 60)</td> <td>1039</td> <td>1027</td> <td>2066</td> </tr> <tr> <td>Gender</td> <td></td> <td></td> <td></td> </tr> <tr> <td>Female</td> <td>4506</td> <td>4605</td> <td>9111</td> </tr> <tr> <td>Male</td> <td>9345</td> <td>9247</td> <td>18592</td> </tr> <tr> <td>Subjects at risk(Co-morbidities)</td> <td>709</td> <td>740</td> <td>1449</td> </tr> <tr> <td>Stable Chronic Heart Disease</td> <td>167</td> <td>155</td> <td>322</td> </tr> <tr> <td>Stable Chronic Lung Disease</td> <td>13</td> <td>7</td> <td>20</td> </tr> <tr> <td>Controlled Diabetic</td> <td>275</td> <td>289</td> <td>564</td> </tr> <tr> <td>Stable Liver Disease</td> <td>2</td> <td>3</td> <td>5</td> </tr> <tr> <td>Severe Obesity</td> <td>18</td> <td>14</td> <td>32</td> </tr> <tr> <td>Other Stable Co-morbid</td> <td>295</td> <td>293</td> <td>588</td> </tr> </tbody> </table> <p>The safety profile of the adolescent age group and the overall population has been found to be same. The common solicited and unsolicited adverse events reported in total population are as under:</p> <p><b>Solicited Local Adverse Events:</b> The most frequently reported solicited local adverse events across all treated subjects in both groups (ZYCOV-D® and Placebo) were pain at injection site: (0.66% and 0.62% after Dose 1; 0.34% and 0.35% after Dose 2 and 0.27% and 0.26% after Dose 3), redness: (0.31% and 0.28% after Dose 1; 0.19% and 0.09% after Dose 2 and 0.17% and 0.09% after Dose 3), swelling: (0.27% and 0.28% after Dose 1; 0.08% and 0.06% after Dose 2 and 0.09% and 0.05% after Dose 3) and itching: 0.08% and 0.14% after Dose 1; 0.05% and 0.07% after Dose 2 and 0.02% and 0.05% after Dose 3). Most of the adverse events were mild or moderate in severity. These events were comparable between ZYCOV-D® and placebo groups.</p> <p><b>Solicited Systemic Adverse Events:</b> The most commonly reported solicited systemic adverse events across all treated subjects in both groups (ZYCOV-D® and Placebo) were headache (0.25% and 0.22% after Dose 1; 0.20% and 0.24% after Dose 2 and 0.16% and 0.17% after Dose 3), fever (0.20% and 0.14% after Dose 1; 0.14% and 0.21% after Dose 2 and 0.13% and 0.10% after Dose 3), muscle pain (0.19% and 0.28% after Dose 1; 0.11% and 0.18% after Dose 2 and 0.11% and 0.09% after Dose 3), and fatigue (0.19% and 0.19% after Dose 1; 0.14% and 0.16% after Dose 2 and 0.09% and 0.13% after Dose 3). Most of the adverse events were mild or moderate in severity. These events were comparable between ZYCOV-D® and placebo groups.</p> <p><b>Unsolicited Adverse Events:</b> Arthralgia, Back pain, Muscle spasms, Myalgia, Musculoskeletal pain, Neck pain, Vertigo, Diarrhoea, Gastritis, Gastroesophageal reflux disease, Nausea, Vomiting, Asthenia, Chills, Eye irritation, Abdominal distension, Abdominal pain, Fatigue, Pain, Pyrexia, Nasopharyngitis, Pain in extremity, Ageusia, Anosmia, Cerebral infarction, Dizziness, Headache, Cough, Dyspnoea, Nasal dryness, Oropharyngeal pain, Rhinorrhoea, Sneezing.</p> <p><b>Serious adverse events:</b> As per interim analysis report, 15 serious adverse events were identified: stroke (02), Death due to Cardiorespiratory arrest with septicemia and alcoholic liver disease (1), Death due to COVID19 (1), Gram negative enteritis (1) and COVID19 (10). None of these serious adverse events was related to IP.</p> <p><b>Adverse Events reported from Phase III Study - 3mg-2dose regimen:</b> A total of 150 adults healthy subjects &gt;18 years of age were enrolled in the Phase III study of 3mg – 2 dose regimen (100 in Vaccine arm and 50 in the Placebo arm). The age group and demographic characteristics of the subjects enrolled in Phase III study are as follows:</p> <table border="1"> <thead> <tr> <th></th> <th>Vaccine (N = 100)</th> <th>Placebo (N = 50)</th> </tr> </thead> <tbody> <tr> <td>Mean Age (years)</td> <td>34.2</td> <td>36.7</td> </tr> <tr> <td>Male</td> <td>68 (68.0%)</td> <td>30 (60.0%)</td> </tr> <tr> <td>Female</td> <td>32 (32.0%)</td> <td>20 (40.0%)</td> </tr> </tbody> </table> <p>The following adverse events were reported during the study: <b>Local site adverse events:</b> Pain (1.9%), Redness (1.4%), Swelling (1.2%), Itching (0.5%) <b>Systemic adverse events:</b> Headache (2.6%), Tiredness / Fatigue (2.1%), Fever (1.1%), Diarrhea (0.5%) and Nausea (0.5%)</p> <p><b>Adverse Events reported from Phase III Study - 3mg-2dose regimen:</b> In our ongoing Phase III clinical trial of 3mg – 2 dose regimen, a total of 3000 healthy subjects &gt;12 years of age have been enrolled. Amongst them 1193 subjects belonged to the adolescent age group (12-17 years). The age group and demographic characteristics of the subjects enrolled in Phase III study are as follows:</p> <table border="1"> <thead> <tr> <th></th> <th>Total (N = 3000)</th> <th>Adults (N = 1807)</th> <th>Adolescents (N = 1193)</th> </tr> </thead> <tbody> <tr> <td>Mean Age (years)</td> <td>27.4</td> <td>35.9</td> <td>14.6</td> </tr> <tr> <td>Male</td> <td>1907 (63.6%)</td> <td>1278 (70.7%)</td> <td>629 (52.7%)</td> </tr> <tr> <td>Female</td> <td>1093 (36.4%)</td> <td>529 (29.3%)</td> <td>564 (47.3%)</td> </tr> </tbody> </table> <p>The safety profile of the adolescent age group and the adult population has been found to be similar. The adverse events reported in total population upto day 112 are as under: <b>Local site adverse events:</b> Pain (1.9%), Redness (0.7%), Swelling (0.4%), Itching (0.3%) <b>Solicited systemic adverse events:</b> Headache (0.4%), Tiredness / Fatigue (0.2%), Fever (0.4%), arthralgia (0.2%), myalgia (0.1%) and vomiting (0.01%). <b>Unsolicited systemic adverse events:</b> Body ache (0.13%), Weakness (0.12%), Headache (0.08%), Cough and Cold (0.08%), Fever (0.07%), Diarrhoea (0.05%), Tiredness (0.03%), COVID-19 (0.06%), Myalgia (0.01%) and Vomiting (0.01%)</p> <p><b>4.9 Overdose</b> Experience of overdose is limited. There is no specific treatment for an overdose with ZYCOV-D®. In the event of an overdose, the individual should be monitored and provided with symptomatic treatment as appropriate.</p> <p><b>5. Pharmacological properties</b> <b>5.1 Mechanism of Action</b> The plasmid construct of ZYCOV-D® carrying the spike-S gene of interest enters host cells, where it remains in the nucleus as an episome; without getting integrated into the host cell DNA. Thus using the host cell's protein translation machinery, the inserted cloned gene in the episome will direct the synthesis of the antigen it encodes. The protein produced by plasmid-transfected cells is likely to be expressed within the cell and folded in its native conformation. Further the signal peptide prompts cells to translocate the protein, usually to the cellular membrane. The antigen is recognized by antigen presenting cells (APCs) and further induces antibodies including neutralizing antibodies and cellular immune response through major histocompatibility complex (MHC).</p> <p><b>5.2 Pharmacodynamic properties</b> <b>Immunogenicity Data 28 days after last dose from Phase II and Phase III Clinical Trials (2mg-3dose regimen with Pharmajet):</b></p> <table border="1"> <thead> <tr> <th colspan="2">Phase II Clinical Trial:</th> </tr> <tr> <th>Parameter</th> <th>Data</th> </tr> </thead> <tbody> <tr> <td>Seroconversion rate based on IgG* (%)</td> <td>91.28%</td> </tr> <tr> <td>Seroconversion rate based on Neutralizing Antibody response<sup>a</sup> (%)</td> <td>88.89%</td> </tr> <tr> <td>GMT based on Neutralizing Antibody response<sup>a</sup></td> <td>131.32 (63.50, 271.58)<sup>‡</sup></td> </tr> <tr> <td>GMFR based on Neutralizing Antibody response<sup>a</sup></td> <td>22.56 (10.57, 48.16)<sup>‡</sup></td> </tr> </tbody> </table> <p><sup>a</sup>by S1 antigen ELISA <sup>b</sup>Wild type virus neutralization assay (PRNT<sub>50</sub>) <sup>‡</sup>data presented as Geometric Mean (95% CI)</p> <table border="1"> <thead> <tr> <th colspan="2">Phase III Clinical Trial:</th> </tr> <tr> <th>Parameter</th> <th>Data</th> </tr> </thead> <tbody> <tr> <td>Seroconversion rate based on IgG* (%)</td> <td>93.33%</td> </tr> <tr> <td>GMT based on IgG<sup>b</sup></td> <td>952.67 (707.9, 1282.0)<sup>‡</sup></td> </tr> <tr> <td>GMFR based on IgG<sup>b</sup></td> <td>136.09 (101.11, 183.1)<sup>‡</sup></td> </tr> </tbody> </table> <p><sup>a</sup>by S1 antigen ELISA <sup>‡</sup>data presented as Geometric Mean (95% CI)</p> <p><b>Interim Efficacy Data from Phase III Clinical Trial (2mg-3dose regimen with Pharmajet):</b> A total of 27703 subjects were enrolled in the Phase III study till interim analysis. The interim primary efficacy analysis was based on the Per-Protocol analysis, which consisted of all participants with negative baseline SARS-CoV-2 status (i.e., negative RT-PCR for SARS-CoV-2) and who had received 3 doses of investigational product. Total of 12350 subjects who had completed 64±3 days in vaccine group and total 12320 subjects who had completed 64±3 days in placebo group were considered for analysis. Out of 81 symptomatic RT-PCR positive COVID-19 cases considered for interim analysis, 61 were in placebo</p>							Phase I: 48 adult subjects						Arm 1, 1 mg (0.1 mL needle and syringe)	Arm 2, 1 mg (0.1 mL Pharmajet)	Arm 3, 2 mg (0.2 mL needle and syringe)	Arm 4, 2 mg (0.2 mL Pharmajet)	Adult subjects (Male)	12	12	12	12	Mean Age (Years)	35.4	31.8	35.1	37.2	Phase II: 1000 subjects						Arm 1, 1 mg (0.1 mL needle and syringe)	Arm 2, 1 mg (0.1 mL Pharmajet)	Arm 3, 2 mg (0.2 mL needle and syringe)	Arm 4, 2 mg (0.2 mL Pharmajet)	N	251	249	250	250	Male	188	186	179	177	Female	63	63	71	73	Adolescent	04	06	06	02	Mean Age (Years)	35.0 ± 11.83	34.2 ± 12.11	35.4 ± 10.46	34.6 ± 10.43	AE Terms	ZyCoV-D n(%)			Placebo n(%)			Dose I (N = 200) n(%)	Dose II (N = 197) n(%)	Dose III (N = 194) n(%)	Dose I (N = 50) n(%)	Dose II (N = 49) n(%)	Dose III (N = 48) n(%)	<b>Solicited Local AEs</b>							Pain at injection site	7 (3.50)	9 (4.57)	6 (3.09)	0 (0.00)	0 (0.00)	0 (0.00)	Redness at injection site	9 (4.50)	10 (5.08)	9 (4.64)	0 (0.00)	0 (0.00)	0 (0.00)	Swelling at injection site	5 (2.50)	6 (3.05)	5 (2.58)	0 (0.00)	0 (0.00)	0 (0.00)	Itching at injection site	1 (0.50)	7 (3.55)	2 (1.03)	0 (0.00)	0 (0.00)	0 (0.00)	Muscle pain	1 (0.50)	0 (0.00)	1 (0.52)	0 (0.00)	0 (0.00)	0 (0.00)	<b>Solicited Systemic AEs</b>							Fatigue	3 (1.50)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	1 (2.08)	Fever	2 (1.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	Headache	3 (1.50)	1 (0.51)	0 (0.00)	1 (2.00)	0 (0.00)	1 (2.08)	Nausea	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	1 (2.08)	<b>Un Solicited Systemic AEs</b>							Covid-19	2 (1.00)	0 (0.00)	2 (1.03)	0 (0.00)	0 (0.00)	1 (2.08)	Nasal Dryness	1 (0.50)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	High Blood Pressure	1 (0.50)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	Arthralgia	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	Body ache	0 (0.00)	3 (1.52)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	Chikungunya virus infection	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	Cough	0 (0.00)	2 (1.02)	1 (0.52)	0 (0.00)	0 (0.00)	0 (0.00)	Pyrexia	0 (0.00)	5 (2.54)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	Headache	0 (0.00)	2 (1.02)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	Myalgia	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	Rhinorrhoea	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	Asthenia	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	Dysuria	0 (0.00)	0 (0.00)	1 (0.52)	0 (0.00)	0 (0.00)	0 (0.00)	Fatigue	0 (0.00)	0 (0.00)	1 (0.52)	0 (0.00)	0 (0.00)	0 (0.00)	Phase III: 27703 subjects (Data at the time of interim analysis), Total sample size: 28216					Vaccine, 2 mg (0.2 mL Pharmajet)	Placebo	Total	Age (in Years) Mean	36.4	36.6	36.5	Age (12-17)	448	487	935	Age (18-60)	12364	12338	24702	Age (above 60)	1039	1027	2066	Gender				Female	4506	4605	9111	Male	9345	9247	18592	Subjects at risk(Co-morbidities)	709	740	1449	Stable Chronic Heart Disease	167	155	322	Stable Chronic Lung Disease	13	7	20	Controlled Diabetic	275	289	564	Stable Liver Disease	2	3	5	Severe Obesity	18	14	32	Other Stable Co-morbid	295	293	588		Vaccine (N = 100)	Placebo (N = 50)	Mean Age (years)	34.2	36.7	Male	68 (68.0%)	30 (60.0%)	Female	32 (32.0%)	20 (40.0%)		Total (N = 3000)	Adults (N = 1807)	Adolescents (N = 1193)	Mean Age (years)	27.4	35.9	14.6	Male	1907 (63.6%)	1278 (70.7%)	629 (52.7%)	Female	1093 (36.4%)	529 (29.3%)	564 (47.3%)	Phase II Clinical Trial:		Parameter	Data	Seroconversion rate based on IgG* (%)	91.28%	Seroconversion rate based on Neutralizing Antibody response <sup>a</sup> (%)	88.89%	GMT based on Neutralizing Antibody response <sup>a</sup>	131.32 (63.50, 271.58) <sup>‡</sup>	GMFR based on Neutralizing Antibody response <sup>a</sup>	22.56 (10.57, 48.16) <sup>‡</sup>	Phase III Clinical Trial:		Parameter	Data	Seroconversion rate based on IgG* (%)	93.33%	GMT based on IgG <sup>b</sup>	952.67 (707.9, 1282.0) <sup>‡</sup>	GMFR based on IgG <sup>b</sup>	136.09 (101.11, 183.1) <sup>‡</sup>
Phase I: 48 adult subjects																																																																																																																																																																																																																																																																																																																																																																																		
	Arm 1, 1 mg (0.1 mL needle and syringe)	Arm 2, 1 mg (0.1 mL Pharmajet)	Arm 3, 2 mg (0.2 mL needle and syringe)	Arm 4, 2 mg (0.2 mL Pharmajet)																																																																																																																																																																																																																																																																																																																																																																														
Adult subjects (Male)	12	12	12	12																																																																																																																																																																																																																																																																																																																																																																														
Mean Age (Years)	35.4	31.8	35.1	37.2																																																																																																																																																																																																																																																																																																																																																																														
Phase II: 1000 subjects																																																																																																																																																																																																																																																																																																																																																																																		
	Arm 1, 1 mg (0.1 mL needle and syringe)	Arm 2, 1 mg (0.1 mL Pharmajet)	Arm 3, 2 mg (0.2 mL needle and syringe)	Arm 4, 2 mg (0.2 mL Pharmajet)																																																																																																																																																																																																																																																																																																																																																																														
N	251	249	250	250																																																																																																																																																																																																																																																																																																																																																																														
Male	188	186	179	177																																																																																																																																																																																																																																																																																																																																																																														
Female	63	63	71	73																																																																																																																																																																																																																																																																																																																																																																														
Adolescent	04	06	06	02																																																																																																																																																																																																																																																																																																																																																																														
Mean Age (Years)	35.0 ± 11.83	34.2 ± 12.11	35.4 ± 10.46	34.6 ± 10.43																																																																																																																																																																																																																																																																																																																																																																														
AE Terms	ZyCoV-D n(%)			Placebo n(%)																																																																																																																																																																																																																																																																																																																																																																														
	Dose I (N = 200) n(%)	Dose II (N = 197) n(%)	Dose III (N = 194) n(%)	Dose I (N = 50) n(%)	Dose II (N = 49) n(%)	Dose III (N = 48) n(%)																																																																																																																																																																																																																																																																																																																																																																												
<b>Solicited Local AEs</b>																																																																																																																																																																																																																																																																																																																																																																																		
Pain at injection site	7 (3.50)	9 (4.57)	6 (3.09)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Redness at injection site	9 (4.50)	10 (5.08)	9 (4.64)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Swelling at injection site	5 (2.50)	6 (3.05)	5 (2.58)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Itching at injection site	1 (0.50)	7 (3.55)	2 (1.03)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Muscle pain	1 (0.50)	0 (0.00)	1 (0.52)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
<b>Solicited Systemic AEs</b>																																																																																																																																																																																																																																																																																																																																																																																		
Fatigue	3 (1.50)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	1 (2.08)																																																																																																																																																																																																																																																																																																																																																																												
Fever	2 (1.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Headache	3 (1.50)	1 (0.51)	0 (0.00)	1 (2.00)	0 (0.00)	1 (2.08)																																																																																																																																																																																																																																																																																																																																																																												
Nausea	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	1 (2.08)																																																																																																																																																																																																																																																																																																																																																																												
<b>Un Solicited Systemic AEs</b>																																																																																																																																																																																																																																																																																																																																																																																		
Covid-19	2 (1.00)	0 (0.00)	2 (1.03)	0 (0.00)	0 (0.00)	1 (2.08)																																																																																																																																																																																																																																																																																																																																																																												
Nasal Dryness	1 (0.50)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
High Blood Pressure	1 (0.50)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Arthralgia	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Body ache	0 (0.00)	3 (1.52)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Chikungunya virus infection	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Cough	0 (0.00)	2 (1.02)	1 (0.52)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Pyrexia	0 (0.00)	5 (2.54)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Headache	0 (0.00)	2 (1.02)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Myalgia	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Rhinorrhoea	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Asthenia	0 (0.00)	1 (0.51)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Dysuria	0 (0.00)	0 (0.00)	1 (0.52)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Fatigue	0 (0.00)	0 (0.00)	1 (0.52)	0 (0.00)	0 (0.00)	0 (0.00)																																																																																																																																																																																																																																																																																																																																																																												
Phase III: 27703 subjects (Data at the time of interim analysis), Total sample size: 28216																																																																																																																																																																																																																																																																																																																																																																																		
	Vaccine, 2 mg (0.2 mL Pharmajet)	Placebo	Total																																																																																																																																																																																																																																																																																																																																																																															
Age (in Years) Mean	36.4	36.6	36.5																																																																																																																																																																																																																																																																																																																																																																															
Age (12-17)	448	487	935																																																																																																																																																																																																																																																																																																																																																																															
Age (18-60)	12364	12338	24702																																																																																																																																																																																																																																																																																																																																																																															
Age (above 60)	1039	1027	2066																																																																																																																																																																																																																																																																																																																																																																															
Gender																																																																																																																																																																																																																																																																																																																																																																																		
Female	4506	4605	9111																																																																																																																																																																																																																																																																																																																																																																															
Male	9345	9247	18592																																																																																																																																																																																																																																																																																																																																																																															
Subjects at risk(Co-morbidities)	709	740	1449																																																																																																																																																																																																																																																																																																																																																																															
Stable Chronic Heart Disease	167	155	322																																																																																																																																																																																																																																																																																																																																																																															
Stable Chronic Lung Disease	13	7	20																																																																																																																																																																																																																																																																																																																																																																															
Controlled Diabetic	275	289	564																																																																																																																																																																																																																																																																																																																																																																															
Stable Liver Disease	2	3	5																																																																																																																																																																																																																																																																																																																																																																															
Severe Obesity	18	14	32																																																																																																																																																																																																																																																																																																																																																																															
Other Stable Co-morbid	295	293	588																																																																																																																																																																																																																																																																																																																																																																															
	Vaccine (N = 100)	Placebo (N = 50)																																																																																																																																																																																																																																																																																																																																																																																
Mean Age (years)	34.2	36.7																																																																																																																																																																																																																																																																																																																																																																																
Male	68 (68.0%)	30 (60.0%)																																																																																																																																																																																																																																																																																																																																																																																
Female	32 (32.0%)	20 (40.0%)																																																																																																																																																																																																																																																																																																																																																																																
	Total (N = 3000)	Adults (N = 1807)	Adolescents (N = 1193)																																																																																																																																																																																																																																																																																																																																																																															
Mean Age (years)	27.4	35.9	14.6																																																																																																																																																																																																																																																																																																																																																																															
Male	1907 (63.6%)	1278 (70.7%)	629 (52.7%)																																																																																																																																																																																																																																																																																																																																																																															
Female	1093 (36.4%)	529 (29.3%)	564 (47.3%)																																																																																																																																																																																																																																																																																																																																																																															
Phase II Clinical Trial:																																																																																																																																																																																																																																																																																																																																																																																		
Parameter	Data																																																																																																																																																																																																																																																																																																																																																																																	
Seroconversion rate based on IgG* (%)	91.28%																																																																																																																																																																																																																																																																																																																																																																																	
Seroconversion rate based on Neutralizing Antibody response <sup>a</sup> (%)	88.89%																																																																																																																																																																																																																																																																																																																																																																																	
GMT based on Neutralizing Antibody response <sup>a</sup>	131.32 (63.50, 271.58) <sup>‡</sup>																																																																																																																																																																																																																																																																																																																																																																																	
GMFR based on Neutralizing Antibody response <sup>a</sup>	22.56 (10.57, 48.16) <sup>‡</sup>																																																																																																																																																																																																																																																																																																																																																																																	
Phase III Clinical Trial:																																																																																																																																																																																																																																																																																																																																																																																		
Parameter	Data																																																																																																																																																																																																																																																																																																																																																																																	
Seroconversion rate based on IgG* (%)	93.33%																																																																																																																																																																																																																																																																																																																																																																																	
GMT based on IgG <sup>b</sup>	952.67 (707.9, 1282.0) <sup>‡</sup>																																																																																																																																																																																																																																																																																																																																																																																	
GMFR based on IgG <sup>b</sup>	136.09 (101.11, 183.1) <sup>‡</sup>																																																																																																																																																																																																																																																																																																																																																																																	